

Informed Consent

Thoracic (Chest) Surgery for Chest Wall Tumor

You have elected to have your pet undergo a surgical procedure called thoracotomy. This is a procedure where the chest is opened surgically to address a problem in the thoracic (chest) cavity. In this case, the surgical goal is to remove a mass, or tumor, of the chest wall. This will require the removal of parts of the ribs, usually a few sections are removed. Generally prognosis with surgery is good, but there are inherent risks with both the thoracotomy and the mass removal. Both will be discussed here. Let us know if you have any questions or concerns.

Thoracotomy

Generally the prognosis with surgery is excellent, complications are rare (10% of cases, or less) but when complications occur, they can be life-threatening. There are many important structures in the chest that require manipulation, treatment of, or dissection near that have the potential to cause complications in chest surgery.

Pneumothorax (air in the chest)- the lungs are built for gas exchange, to get oxygen in to the blood, but air in the chest, outside of the lungs, does not allow that gas exchange due to pressure changes. As soon as the chest is opened to room air during surgery, your pet loses the negative pressure the diaphragm obtains in the chest used to pull air into the lungs. This is ok when it is planned, as in during surgery, because we can provide pressure by ventilating your pet during the procedure. If there is a leakage of air into the chest (from the environment or from a defect or damage in a lung or airway) after surgery, this can be life threatening. Most often, we recommend a chest tube be placed after surgery to be sure “negative pressure” is obtained for a few days to ensure the gas spaces are appropriately sealed.

Hemothorax (blood in the chest)- if there is continued hemorrhage after surgery, it collects in the chest, outside of the lungs. If there is sufficient volume of blood, it may not only raise concerns about low blood volume, but also the lungs may not have room to expand to allow breathing. This could be life-threatening.

Pyothorax (pus in the chest)- this could be due to the underlying process or could be secondary to surgery. Additional treatments (chest tube) or procedure may be recommended.

Nerve damage- specifically the phrenic nerve could be injured during surgery or dissection. The phrenic nerve is in the chest and runs along the heart. There are two (right and left)

and they are responsible for helping the diaphragm function (contract to allow breathing). Damage to this structure could make it harder (permanently or temporarily, depending on the amount of damage) for your pet to breathe after surgery.

Cardiac arrhythmias- these are disturbances in the heart rhythm, and may be due to irritation of the heart or damage to heart muscle. Usually they are temporary and may or may not need medication.

Complications with chest tubes- the purpose is to monitor and evacuate the thoracic space. However, they have their own set of issues: their presence or placement can damage structures in the chest, leading to bleeding or pneumothorax (see above), heart muscle damage, heart rhythm disturbances (arrhythmias), or nerve damage, they could become displaced or leak (and lead to pneumothorax), or they could block or plug. Proper care, monitoring, and maintenance are key.

More minor complications of thoracic surgery include wound infection, wound dehiscence (opening), seroma (collection of fluid under the skin), subcutaneous emphysema (collection of air under the skin).

Rare complications include infection of the bone or ribs (osteomyelitis, a serious infection), fractured or displaced ribs/sternum, and lameness in the front limb(s) due to stretching of nerves in the armpits during surgery (usually temporary).

Chest Wall Tumor

Generally, perioperative risks depend on the size and nature of the tumor. However, here are the most common things that may occur (in about 30% of cases):

Failure to close the chest wall: in some cases a synthetic mesh or suture can be used to create a barrier between the chest and the skin, but reconstructive procedures may be needed

Wound complications: infection, dehiscence (opening of the wound), seroma (fluid collection/swelling under the skin)

Tumor recurrence- the tumor may grow back or spread to another area of the body

Diaphragmatic hernia- in some cases, the diaphragm is affected, part of it must be removed or moved and if the suture line fails, that could lead to herniation